

Correction: Galbanic acid of *Ferula assa-foetida* L, as a regulator of the AMPK pathway in reduction of lipid accumulation in HepG2 cells

Hadis Musavi^{1,2}^{ID}, Hajar Shokri-Afra³^{ID}, Soleiman Mahjoub^{2,4*}^{ID}, Abbas Khonakdar-Tarsi^{5,6*}^{ID}, Abouzar Bagheri^{5,6}^{ID}, Zahra Memariani⁷^{ID}

¹Student Research Committee, Babol University of Medical Sciences, Babol, Iran

²Department of Clinical Biochemistry, School of Medicine, Babol University of Medical Sciences, Babol, Iran

³Gut and Liver Research Center, Non-communicable Disease Institute, Mazandaran University of Medical Sciences, Sari, Iran

⁴Cellular and Molecular Biology Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran

⁵Molecular and Cell Biology Research Center, Faculty of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

⁶Department of Clinical Biochemistry and Genetics, Faculty of Medicine, Mazandaran University of Medical Sciences, Sari, Iran

⁷Traditional Medicine and History of Medical Sciences Research Center, Health Research Institute, Babol University of Medical Sciences, Babol, Iran

*Correspondence to

Soleiman Mahjoub, Email: smahjoub20@gmail.com, s.mahjoub@mubabol.ac.ir and Abbas Khonakdar-Tarsi, Email: khonakdarab@gmail.com

Received 17 May 2024

Accepted 7 Aug. 2024

Published online 6 Oct. 2024

Correction

This document corrects the article titled “Galbanic acid of *Ferula assa-foetida* L. as a regulator of the AMPK pathway in the reduction of lipid accumulation in HepG2 cells,” published in 2023, Volume 09, Issue 02, DOI: 10.34172/ipp.2023.39479. The original version of the article contained an error in the uploading of image B in Figure 2. This issue has now been rectified in the PDF version of the article. The following figure represents the corrected version of Figure 2.




Figure 2. Effect of Gal/RSV on HepG2 cells. (A) Dose-dependent effect of Gal on HepG2 cell survival. Cells with the desired concentrations of Gal were treated for 24 hours and then MTT was performed. (B) effect of Gal/RSV on intracellular total lipid content of HepG2 cells with 0.4 mM Pal alone, with 300 µM Gal, and with 50 µM RSV were treated with 0.4 mM palmitate for 24 hours. (C) shows the semi-quantitative results of ORO staining of the effect of Gal/RSV on the induction of lipid accumulation in HepG2 cells. Results are mean ± SD from three independent experiments. * $P < 0.05$, and ** $P < 0.01$ compared with the untreated control.

Citation: Musavi H, Shokri-Afra H, Mahjoub S, Khonakdar-Tarsi A, Bagheri A, Memariani Z. Correction: Galbanic acid of *Ferula assa-foetida* L, as a regulator of the AMPK pathway in reduction of lipid accumulation in HepG2 cells. Immunopathol Persa. 2025;x(x):e39479. DOI:10.34172/ipp.2025.40672.

